

The invariant field of the adjoint action of the unitriangular group in the nilradical of a parabolic subalgebra

V. V. Sevostyanova*

ABSTRACT. In the present paper the field of invariants of the adjoint action of the unitriangular group in the nilradical of any parabolic subalgebra is described.

Consider the general linear group $GL(n, K)$ defined over an algebraically closed field K of characteristic 0. Let B (N , respectively) be its Borel (maximal unipotent, respectively) subgroup, which consists of triangular matrices with nonzero (unit, respectively) elements on the diagonal. We fix a parabolic subgroup P that contains B . Denote by $\mathfrak{p}$, $\mathfrak{b}$ and $\mathfrak{n}$ the Lie subalgebras in $\mathfrak{gl}(n, K)$ that correspond to P , B and N respectively. We represent $\mathfrak{p} = \mathfrak{r} \oplus \mathfrak{m}$ as the direct sum of the nilradical $\mathfrak{m}$ and a block diagonal subalgebra $\mathfrak{r}$ with sizes of blocks $(n_1, \dots, n_s)$. The subalgebra $\mathfrak{m}$ is invariant relative to the adjoint action of the group P , therefore, $\mathfrak{m}$ is invariant relative to the action of the subgroups B and N . We extend this action to the representation in the algebra $K[\mathfrak{m}]$ and in the field $K(\mathfrak{m})$. The subalgebra $\mathfrak{m}$ contains a Zariski-open P -orbit, which is called the *Richardson orbit* (see [R]). Consequently, the algebra of invariants $K[\mathfrak{m}]^P$ coincides with K . The question concerning the structure of the algebra of invariants $K[\mathfrak{m}]^N$ and $K[\mathfrak{m}]^B$ remains open and seems to be a considerable challenge. In the special case $P = B$, the answer is following: the algebra of invariants $K[\mathfrak{m}]^N$ is the polynomial algebra $K[x_{12}, x_{23}, \dots, x_{n-1,n}]$. The present paper is a continuation of a paper [PS], in which a conjecture on the structure of the field of invariants $K(\mathfrak{m})^N$ was stated. This conjecture is proved in the present paper. Theorem 1.8 provides a complete description of the field of invariants $K(\mathfrak{m})^N$ for any parabolic subalgebra. Another result of the paper is Theorem 1.7, in which canonical representation of N -orbits in general position are indicated.

*This research was partially supported by the RFBR (project 08-01-00151-a)

1. The main definitions and results

Every positive root γ in $\mathfrak{gl}(n, K)$ has the form (see [GG]) $\gamma = \varepsilon_i - \varepsilon_j$, $1 \leq i < j \leq n$. We identify a root γ with the pair (i, j) and the set of the positive roots Δ^+ with the set of pairs (i, j) , $i < j$. The system of positive roots $\Delta_{\mathfrak{r}}^+$ of the reductive subalgebra $\mathfrak{r}$ is a subsystem in Δ^+ .

Let $\{E_{i,j} : i < j\}$ be the standard basis in $\mathfrak{n}$. By E_{γ} denote the basis element $E_{i,j}$, where $\gamma = (i, j)$.

We define a relation in Δ^+ such that $\gamma' \succ \gamma$ whenever $\gamma' - \gamma \in \Delta_{\mathfrak{r}}^+$. If $\gamma \prec \gamma'$ or $\gamma \succ \gamma'$, then the roots γ and γ' *comparable*. Denote by M the set of $\gamma \in \Delta^+$ such that $E_{\gamma} \in \mathfrak{m}$. We identify the algebra $K[\mathfrak{m}]$ with the polynomial algebra in the variables $x_{i,j}$, $(i, j) \in M$.

Definition 1.1. A subset S in M is called a *base* if the elements in S are not pairwise comparable and for any $\gamma \in M \setminus S$ there exists $\xi \in S$ such that $\gamma \succ \xi$.

Definition 1.2. Let A be a subset in S . We say that γ is a *minimal element* in A if there is no $\xi \in A$ such that $\gamma \succ \xi$.

Note that M has a unique base S , which can be constructed in the following way. We form the set S_1 of minimal elements in M . By definition, $S_1 \subset S$. Then we form a set M_1 , which is obtained from M by deleting S_1 and all elements

$$\{\gamma \in M : \exists \xi \in S_1, \gamma \succ \xi\}.$$

The set of minimal elements S_2 in M_1 is also contained in S , and so on. Continuing the process, we get the base S .

Definition 1.3. An ordered set of positive roots $\{\gamma_1, \dots, \gamma_s\}$ is called a *chain* if $\gamma_1 = (a_1, a_2)$, $\gamma_2 = (a_2, a_3)$, $\gamma_3 = (a_3, a_4)$, and so on. The number s is called the *length of a chain*.

Definition 1.4. We say that two roots $\xi, \xi' \in S$ form an *admissible pair* $q = (\xi, \xi')$ if there exists $\alpha_q \in \Delta_{\mathfrak{r}}^+$ such that the ordered set of roots $\{\xi, \alpha_q, \xi'\}$ is a chain. Note that the root α_q is uniquely determined by q .

We form the set $Q := Q(\mathfrak{p})$ that consists of admissible pairs of roots in S . For every admissible pair $q = (\xi, \xi')$ we construct a positive root $\phi_q = \alpha_q + \xi'$. Consider the subset $\Phi = \{\phi_q : q \in Q\}$.

Using the given parabolic subgroup, we construct a diagram, which is a square matrix in which the roots from S are marked by the symbol $\otimes$ and the roots from Φ are labeled by the symbol $\times$. The other entries in the diagram are empty.

Example 1. Below a diagram for a parabolic subalgebra with sizes of its diagonal blocks $(1, 3, 2, 1, 3, 2, 2)$ is given.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	
1	1	⊗													1
2		1			×	×				⊗					2
3			1			⊗									3
4				1	⊗										4
5					1	×		⊗							5
6						1	⊗								6
7							1	⊗							7
8								1			×	×			8
9									1		×	⊗			9
10										1	⊗				10
11											1	×	⊗		11
12												1	⊗		12
13													1		13
14														1	14

Diagram 1.

Consider the formal matrix $\mathbb{X}$ in which the variables $x_{i,j}$ occupy the positions $(i,j) \in M$ and the other entries are equal to zero. For any root $\gamma = (a,b) \in M$ we denote by S_γ the set of $\xi = (i,j) \in S$ such that $i > a$ and $j < b$. Let $S_\gamma = \{(i_1, j_1), \dots, (i_k, j_k)\}$. Denote by M_γ a minor M_I^J of the matrix $\mathbb{X}$ with ordered systems of rows I and columns J , where

$$I = \text{ord}\{a, i_1, \dots, i_k\}, \quad J = \text{ord}\{j_1, \dots, j_k, b\}.$$

For every admissible pair $q = (\xi, \xi')$, we construct the polynomial

$$L_q = \sum_{\substack{\alpha_1, \alpha_2 \in \Delta_+^\perp \cup \{0\} \\ \alpha_1 + \alpha_2 = \alpha_q}} M_{\xi + \alpha_1} M_{\alpha_2 + \xi'}. \quad (1)$$

Theorem 1.5 [PS]. *For an arbitrary parabolic subalgebra, the system of polynomials*

$$\{M_\xi, \xi \in S, L_\varphi, \varphi \in \Phi, \}$$

is contained in $K[\mathfrak{m}]^N$ and is algebraically independent over K .

Denote by $\mathcal{Y}$ the subset in $\mathfrak{m}$ that consists of matrices of the form

$$\sum_{\xi \in S} c_\xi E_\xi + \sum_{\varphi \in \Phi} c'_\varphi E_\varphi,$$

where $c_\xi \neq 0$ and $c'_\varphi \neq 0$.

Definition 1.6. The matrices from $\mathcal{Y}$ are said to be *canonical*.

In the fourth section, we prove the following two theorems.

Theorem 1.7. *There exists a nonempty Zariski-open subset $U \subset \mathfrak{m}$ such that the N -orbit of any $x \in U$ intersects $\mathcal{Y}$ at a unique point.*

Theorem 1.8. *The field of invariants $K(\mathfrak{m})^N$ is the field of rational functions of M_ξ , $\xi \in S$, and L_φ , $\varphi \in \Phi$.*

The following result is a consequence of Theorem 1.8.

Theorem 1.9. *The maximal dimension of an N -orbit in $\mathfrak{m}$ is equal to $\dim \mathfrak{m} - |S| - |\Phi|$.*

We introduce on the set of roots $S \cup \Phi$ an order relation for which

- 1) $\xi < \varphi$ for any $\xi \in S$ and $\varphi \in \Phi$;
- 2) for other pairs of roots from $S \cup \Phi$, the relation $<$ means that $(a, b) < (c, d)$ if $c < a$ or $c = a$ if $b < d$.

Consider the roots from the set $S \cup \Phi$: $S = \{\xi_1, \dots, \xi_p\}$, $\Phi = \{\varphi_1, \dots, \varphi_q\}$. Let the mapping $\varpi : \mathfrak{m} \rightarrow K^{p+q}$ be given by

$$x \mapsto (M_{\xi_1}(x), \dots, M_{\xi_p}(x), L_{\varphi_1}(x), \dots, L_{\varphi_q}(x)).$$

Denote $(c_{\xi_1}, \dots, c_{\xi_p}, c_{\varphi_1}, \dots, c_{\varphi_q}) \in \varpi(\mathfrak{m})$. Let $\alpha \in S$. By $\widehat{c}_\alpha$ denote the number $\prod c_\beta$, where the product proceeds over all roots $\beta \in S$ such that $\beta < \alpha$ and β is the greatest root in the sense of the order $<$. Let U be a Zariski-open set in the condition of Theorem 1.7. We have a canonical representative of the N -orbit of any matrix from U .

Corollary 1.10. *Let $x \in U$ and $\varpi(x) = (c_{\xi_1}, \dots, c_{\xi_p}, c_{\varphi_1}, \dots, c_{\varphi_q})$. Then the canonical representative of the N -orbit of the element x is the following element*

$$\sum_{i=1}^p \frac{c_{\xi_i}}{\widehat{c}_{\xi_i}} E_{\xi_i} + \sum_{j=1}^q \frac{c_{\varphi_j}}{c_{\gamma_j} \widehat{c}_{\gamma'_j}} E_{\varphi_j},$$

where $\varphi_j \in \Phi$ corresponds to the admissible pair (γ_j, γ'_j) .

2. Lemmas concerning the structure of the root systems S and Φ .

In this section we prove several lemmas. As above, let $(r_1, \dots, r_s)$ be the sizes of blocks in the reductive subalgebra $\mathfrak{r}$. We denote $R_k = \sum_{t=1}^k r_t$.

Lemma 2.1. *For any root $(i, j) \in M \setminus S$, there exists a number $\tilde{i} > i$ such that $(\tilde{i}, j) \in S$ or a number $\tilde{j} < j$ for which $(i, \tilde{j}) \in S$.*

Remark. In other words, the base S is so structured that for any cell (i, j) of the matrix in $\mathbf{m}$, there exists an element of the base S that lies in the i th row and/or in the j th column.

PROOF. By definition of the base S , for arbitrary $\gamma = (i, j)$ in $M \setminus S$ there exists a root $\xi = (k, m) \in S$ such that $\gamma - \xi \in \Delta^+$, i.e.,

$$\gamma - \xi = \varepsilon_i - \varepsilon_j - (\varepsilon_k - \varepsilon_m) \in \Delta^+.$$

The last inclusion holds if $i = k$ or $j = m$. $\square$

Lemma 2.2. *Suppose that a root $(i, j) \in S$ is such that $i \neq R_k$ for any k . Then there exists a number $\tilde{j} < j$ such that $(i + 1, \tilde{j}) \in S$.*

PROOF. Consider the root $(i + 1, j) \in M$; it does not lie in S and beneath it there are no roots from S , because otherwise there are two roots from S in one column. Assume that for any $\tilde{j} < j$ we have $(i + 1, \tilde{j}) \notin S$. Then for the root $(i + 1, j)$ there is no $\xi \in S$ for which

$$(i + 1, j) - \xi \in \Delta^+,$$

which contradicts the definition of the base. $\square$

Corollary. *Suppose $(i, j) \in S$ is such that for some number k , we have*

$$R_{k-1} < i < R_k.$$

Then for any $\tilde{i}$ such that $i < \tilde{i} \leq R_k$, there exists a number $\tilde{j} < j$ for which $(\tilde{i}, \tilde{j}) \in S$.

PROOF proceeds by induction on the number $\tilde{i}$. $\square$

The following lemma is proved similarly.

Lemma 2.3. *Suppose a root $(i, j) \in S$ is such that $j \neq R_k + 1$ for any k . Then there exists a number $\tilde{i} > i$ such that the root $(\tilde{i}, j - 1) \in S$.*

Corollary. *Assume that $(i, j) \in S$ is such that for some number k , we have*

$$R_k + 1 < j \leq R_{k+1}.$$

Then for any $\tilde{j}$ satisfying $R_k < \tilde{j} < j$, there exists a number $\tilde{i} > i$ such that $(\tilde{i}, \tilde{j}) \in S$.

Lemma 2.4. *Let a root $(i, j) \in S$ be such that there is no root $(\tilde{i}, \tilde{j}) \in S$ with $i > \tilde{i}$ and $j < \tilde{j}$. Assume that for some a and b , the following conditions are satisfied: $R_a < i \leq R_{a+1}$, $R_{b-1} < j \leq R_b$. Then $i = 1 + R_a$ or $j = R_b$.*

For example, in Diagram 1 the roots $(1, 2)$, $(2, 10)$, $(9, 12)$ and $(11, 14)$ satisfy the assumptions of Lemma 2.4.

PROOF. Assume the contrary, i.e., $i > R_a + 1$ and $j < R_b$.

Consider the root $(i - 1, j + 1) \in M$; from the conditions of the lemma, it follows that it does not lie in S . Then there exists $\xi \in S$ for which

$$(i - 1, j + 1) - \xi \in \Delta^+.$$

Lemma 2.1 implies that $\xi = (i - 1, k)$, where $k < j + 1$, or $\xi = (m, j + 1)$, where $i - 1 < m$. In the first case, by the corollary of Lemma 2.2, for any number k such that $R_{b-1} < k < j + 1$, there exists a root $(i_k, k) \in S$, where $i_k > i - 1$. Then $\xi \notin S$, because otherwise in the k th column there are two roots from S , which contradicts the minimality of the base. The second case is also impossible by the consequence of Lemma 2.3. Therefore, such a $\xi \in S$ does not exist and $(i - 1, j + 1) \in S$, a contradiction. $\square$

Lemma 2.5. *Suppose the following r roots lie in S :*

$$(i_1, j - r + 1), (i_2, j - r + 2), \dots, (i_r, j) = (i, j), \quad (2)$$

where $i_1 > i_2 > \dots > i_r$; moreover, there is no root $(i_0, j - r) \in S$ such that $i_0 > i_1$. Then there are precisely r roots in S of the form

$$(i + r - 1, j_1), (i + r - 2, j_2), \dots, (i, j_r) = (i, j), \quad (3)$$

where $j_1 < j_2 < \dots < j_r$.

The converse is also true; namely, if r roots of the form (3) lie in S and there is no root $(i - r, j_0) \in S$ with $j_0 < j_1$, then there are precisely r roots in S of the form (2).

PROOF. The corollary to Lemma 2.2 asserts that such roots exist. Let the number of them be equal to t :

$$(i + t - 1, j_1), (i + t - 2, j_2), \dots, (i, j_t) = (i, j), \quad j_k < j_{k+1}. \quad (4)$$

The first root $(i_1, j - r + 1)$ on the list (2) is minimal (in the sense of the relation $\succ$), and the value of j_1 is the least possible. Then $i_1 + 1 = j - r + 1$, consequently, $i_1 = j - r$. Similarly, the first root in (4) is also minimal; therefore, we have $i + t - 1 + 1 = j_1$, i.e., $j_1 = i + t$. Since the root $\xi =$

$(i, j) \in S$, it follows that a square minor M_ξ corresponds to it. We find its size: $i_1 - i + 1 = j - j_1 + 1$, whence we have $j - r - i = j - (i + t)$ and $r = t$. $\square$

Definition 2.6. Any root $(i, j) \in M$ satisfying the conditions

$$R_{k-1} < i \leq R_k \text{ and } R_k < j \leq n,$$

is called a *root lying to the right of the k th block* in $\mathfrak{r}$.

Lemma 2.7. *Let roots*

$$(i, j_1), (i, j_2), \dots, (i, j_a) \in S \cup \Phi,$$

where $j_1 < j_2 < \dots < j_a$ and $a > 1$, and let other roots from $S \cup \Phi$ do not lie in the i th row. Then

- 1) if $R_{l-1} < i \leq R_l$, then $j_1 = R_l + 1$;
- 2) if $j_1 \leq R_k < j_b$ for some number $b \leq a$, then $r_k < b$.

PROOF.

- 1). We prove the first item of the lemma. Let $R_{l-1} < i \leq R_l$ and $j_1 \neq R_l + 1$. Since the root $(i, j_1) \in \Phi$, there exist roots $(\tilde{i}, j_1)$, $\tilde{i} > i$, and (c, i) that lie in S . Next, since $j_1 \neq R_l + 1$, by Lemma 2.3 there is a root $(\hat{i}, j_1 - 1) \in S$. Then the pair of roots $((c, i), (\hat{i}, j_1 - 1))$ is admissible. Hence we have $(i, j_1 - 1) \in \Phi$, a contradiction. Thus $j_1 = R_l + 1$.
- 2). Let we have $R_{m-1} < j_b \leq R_m$ for some number m . Show that $r_k < b$ for any k satisfying the condition $l < k < m$. We prove this by induction on k . For the $(l+1)$ th block in $\mathfrak{r}$ the statement is obvious. Suppose the sizes of blocks in $\mathfrak{r}$ the numbers of which is equal to $l+1, l+2, \dots, k-1$ are less than b . We show that $r_k < b$.

Suppose the contrary. Let $r_k > b$; then for any number j , where $R_{k-1} < j \leq R_k$, a root $(a_j, j) \in S$ exists for some $a_j \geq i$. Thus, there are r_k roots from S of the form

$$(i_1, R_{k-1} + 1), (i_2, R_{k-1} + 2), \dots, (i_{r_k}, R_k)$$

for some $i_1 > i_2 > \dots > i_{r_k}$. Lemma 2.5 implies that there exist r_k roots in S that lie to the right of a certain block in $\mathfrak{r}$. Then the size of this block is no less than r_k and thus is greater than b . Since its number is less than k , we obtain a contradiction with the inductive assumption. Consequently, for any number k such that $j_1 \leq R_k < j_b$, we have $r_k < b$. $\square$

3. The root system T and the principal minors

Denote by $w_{i,j} : \mathfrak{m} \rightarrow K$ the coordinate function that maps a matrix $x \in \mathfrak{m}$ to the number that stands at the intersection of the i th row and j th column.

Consider $x \in \mathfrak{m}$. Let the corresponding block diagonal subalgebra $\mathfrak{r}$ consists of blocks of sizes $(r_1, \dots, r_s)$. Then $n = r_1 + \dots + r_s$. We fix the size of the last block: $r_s = r$. As above, $R_k = \sum_{t=1}^k r_t$.

To prove Theorem 1.7, we need to consider two cases, where there is a root from S in the n th column and there are no roots from S in the n th column. To prove the theorem in the first case, we introduce the notion of principal minor (Definition 3.3) and define root systems Ψ and T (notation 3.1 and 3.6, respectively). Throughout the present section, we assume that a parabolic subalgebra $\mathfrak{p}$ is such that there is a root from S in the n th column. We denote it by $(\tilde{m}, n)$. We take a number $m \leq \tilde{m}$ such that $(m, n) \in S \cup \Phi$ and if $i < m$, then $(i, n) \notin S \cup \Phi$. By Lemma 2.5 and the corollary to Lemma 2.2, there exists precisely r roots in S of the form

$$(\tilde{m} + r - 1, l_1), (\tilde{m} + r - 2, l_2), \dots, (\tilde{m} + 1, l_{r-1}), (\tilde{m}, l_r) = (\tilde{m}, n)$$

for which $l_1 < l_2 < \dots < l_r = n$.

We show graphically the part of the diagram for $\mathfrak{m}$ that contains these roots:

	l_1	l_2	l_{r-1}	$l_r = n$		
1	×	...	×	...	×	m
...						
1	×	...	×	...	×	$\tilde{m} - 1$
1	×	...	×	...	×	$\tilde{m}$
1		...		...	⊗	
...						
1		...	⊗	...		$\tilde{m} + r - 2$
1	⊗					$\tilde{m} + r - 1$

Consider all possible chains of roots from $S \cup \Phi$ that contain the roots (i, j) , where $m \leq i < m + r$ and $j \in \{l_1, l_2, \dots, l_r\}$. Among these chains, we chose those satisfying the following condition: if (a, b) is a root of such chain and $R_k < a \leq R_{k+1}$ for some $k \geq 1$, then

$$a \leq r + R_k. \quad (5)$$

Notation 3.1. The set of all roots that occur in such chains is denoted by Ψ .

Obviously, the length of any root chain from $S \cup \Phi$ is less than the number of blocks in $\mathfrak{r}$. With Example 1, we show the arrangement of roots from Ψ . We mark them by the symbol $\boxtimes$.




Diagram 2

We recall that s is the number of blocks of the reductive subalgebra $\mathfrak{r}$. The size of the last block in $\mathfrak{r}$ is equal to r . Obviously, for some number $\tilde{s} < s$ we have $m = R_{\tilde{s}} + 1$. Lemma 2.7 implies that if $\tilde{s} + 1 < t < s$, then the size of the t th block in $\mathfrak{r}$ is less than the size of the last block r . For this reason, all roots (i, j) lying in $S \cup \Phi$ such that $i > R_{\tilde{s}+1}$ are contained in the root system Ψ .

Example 2. We give one more example of the diagram of a parabolic subalgebra with sizes of diagonal blocks $(2, 2, 1, 3, 2, 1, 3)$. Here $\tilde{s} = 3$ and $m = R_3 + 1 = 6$.




Diagram 3

Denote $\tilde{r}_k = \min(r, r_k)$. Then from the definition of the set Ψ it follows that if $(i, j) \in \Psi$, then for $k > 1$ we have

$$R_{k-1} < i \leq \tilde{r}_k + R_{k-1}.$$

Lemma 3.2. *Let $k > 1$. The following statement are valid:*

1. *There are $\tilde{r}_k$ roots from $S \cup \Phi$ in the row with number $R_{k-1} + 1$.*
2. *The first (according to the number of a column) $\tilde{r}_k$ roots from $S \cup \Phi$ in the row with number $R_{k-1} + 1$ lie in Ψ .*
3. *All roots from Ψ that lie to the right of the k th block in $\mathfrak{r}$ are contained in the rows with numbers $R_{k-1} + 1, R_{k-1} + 2, \dots, R_{k-1} + \tilde{r}_k$ and in the same columns as the roots from Ψ in the row with number $R_{k-1} + 1$.*

PROOF.

1. (a) First we prove that at least $\tilde{r}_k$ roots from S lie to the right of the k th block.

The proof proceeds by contradiction. assume that to the right of the k th block, there are less than $\tilde{r}_k$ roots from S . Then there are no roots from S in the row with the number $R_{k-1} + 1$. Otherwise to the right of the k th block there are $r_k \geq \tilde{r}_k$ roots from S . By Lemma 2.1, for each root $(R_{k-1} + 1, j) \in M$ lying in the row with number $R_{k-1} + 1$, a root $\xi = (i, j) \in S$ with $i > R_{k-1} + 1$ exists.

Obviously, $r = r_s \geq \tilde{r}_k$. Let l be the least number for which $r_l \geq \tilde{r}_k$, $l > k$. Since $r_s = r \geq \tilde{r}_k$ and $s > k$, we have $l \leq s$.

Next, in each of the columns with numbers

$$R_{l-1} + 1, R_{l-1} + 2, \dots, R_{l-1} + r_l = R_l$$

there is a root $(a_i, R_{l-1} + i) \in S$, where $i = 1, \dots, r_l$. Lemma 2.3 implies that $a_1 > a_2 > \dots > a_{r_l}$. Then, by Lemma 2.5, there exists a number $k \leq p < l$ such that there are r_l roots from S to the right of the p th block in $\mathfrak{r}$. Then $r_p \geq r_l \geq \tilde{r}_k$. Since l was chosen to be minimal, we obtain $l = k$. But this contradicts the assumption that to the right of the k th block there are less than $\tilde{r}_k$ roots from S . Thus, to the right of the k th block there are $\tilde{r}_k$ roots from S . Part (a) is proved.

- (b) We complete the proof of statement 1. Let the size of the k th block in $\mathfrak{r}$ be greater than 1. For $r_k = 1$ the proof is trivial, because $(R_k, R_k + 1) \in S$ for any number k .

The corollary to Lemma 2.3 states that the roots from S lying to the right of the k th block in $\mathfrak{r}$ have the form

$$\xi_1 = (R_k, j_1), \xi_2 = (R_k - 1, j_2), \dots, \xi_{\tilde{r}_k} = (R_k - \tilde{r}_k + 1, j_{\tilde{r}_k}), \dots,$$

where $j_1 = R_k + 1$ and $j_1 < j_2 < \dots < j_{\tilde{r}_k}$. We show that all the roots of the form $(R_{k-1} + 1, j_a)$, where $1 \leq a \leq \tilde{r}_k$, lie in $S \cup \Phi$.

Consider the root $\eta = (R_{k-1}, R_{k-1} + 1)$; since $k > 1$, this root obviously lies in S . Denote

$$I = \begin{cases} 1, 2, \dots, \tilde{r}_k - 1, \text{ если } \tilde{r}_k = r_k; \\ 1, 2, \dots, \tilde{r}_k, \text{ если } \tilde{r}_k < r_k. \end{cases}$$

Every root $\gamma_a = (R_{k-1} + 1, R_k - a + 1)$, where $a \in I$, lies in $\Delta_{\mathfrak{r}}^+$. Consequently, all pairs of roots (η, ξ_a) , $a \in I$, are admissible. Denote

$$\zeta_a = \begin{cases} \gamma_a + \xi_a = (R_{k-1} + 1, j_a), \text{ если } \tilde{r}_k < r_k, \\ \zeta_{\tilde{r}_k} = \xi_{\tilde{r}_k}, \text{ если } \tilde{r}_k = r_k. \end{cases} \quad (6)$$

Then the roots ζ_a are in Φ for any $a \in I$, which proves the first statement of the lemma.

We prove statement 2 and 3 for numbers k that do not exceed $\tilde{s} + 1$. For remaining k , there is nothing to prove.

2. The proof of the second statement proceeds by induction on the number k , starting with the greatest one. For $k = \tilde{s} + 1$ the statement is obvious. Suppose statement 2 is valid for roots from Ψ that lie to the right of the blocks in $\mathfrak{r}$ with numbers $k + 1, k + 2, \dots, \tilde{s} + 1$. we show this for the block with number k ; namely, we prove that for any $1 \leq a \leq \tilde{r}_k$, the root ζ_a , defined in (6), lie in the root system Ψ .

Lemma 2.5 implies that there exist roots in S of the form

$$(i_1, j_a - a + 1), (i_2, j_a - a + 2), \dots, (i_a, j_a) = (R_k - a + 1, j_a)$$

for some $i_1 > i_2 > \dots > i_a = R_k - a + 1$. For the first root on the list, we have $i_1 = j_a - a = R_b$ for some number $b \geq k$.

Denote $\xi = (R_{b+1}, R_{b+1} + 1)$. It is easy to see that $\xi \in S$. If $j_a < R_{b+1}$, then $(j_a, R_{b+1}) \in \Delta_{\mathfrak{r}}^+$; then the pair (ξ_a, ξ) is admissible. Consequently, the root $(j_a, R_{b+1} + 1)$ lies in Φ . If $j_a = R_{b+1}$, then $(j_a, R_{b+1} + 1)$ lies in the base S , i.e., in either case ($j_a < R_{b+1}$ and $j_a = R_{b+1}$), the roots $(j_a, R_{b+1} + 1) \in S \cup \Phi$. It remains to prove that $(j_a, R_{b+1} + 1) \in \Psi$. Consider the chain

$$(j_a, R_{b+1} + 1), (R_{b+1} + 1, R_{b+2} + 1), (R_{b+2} + 1, R_{b+3} + 1), \dots, (R_{\tilde{s}} + 1, R_{\tilde{s}+1} + 1),$$

all of its roots lie in $S \cup \Phi$. Since

$$j_a = a + R_b \leq \tilde{r}_k + R_b \leq r + R_b,$$

the pair $(j_a, R_{b+1} + 1)$ satisfies (5). Consequently $(j_a, R_{b+1} + 1) \in \Psi$. Hence the root ζ_a also lies in Ψ .

3. We prove statement 3. Consider a root (i, j) , where $R_{k-1} + 1 < i \leq R_{k-1} + \tilde{r}_k$, $j_{a-1} < j < j_a$ for some $a \in \{2, \dots, \tilde{r}_k\}$. Suppose $(i, j) \in S \cup \Psi$. Therefore, there is a root $(\tilde{i}, j) \in S$, $\tilde{i} > i$. Then, similarly to item 1 and 2 of the lemma, we obtain $(R_{k-1} + 1, j) \in \Psi$, a contradiction with $j \neq j_a$ for any a . Thus, $(i, j) \notin \Psi$.

It remains to show that there are no roots from Ψ that lie in the rectangle $R_{k-1} < i \leq R_{k-1} + \tilde{r}_k$, $j_{\tilde{r}_k} < j \leq n$. Assume the contrary. Let $(i, j) \in \Psi$. Then $(i, j) \in S \cup \Phi$ and there exists a number $a > \tilde{r}_k$ such that $\xi_a = (R_k - a + 1, j) \in S$, where $R_k - a + 1 \geq i$.

Note that if $a > \tilde{r}_k$, then $r_k > \tilde{r}_k$, whence $\tilde{r}_k = r$ and $a > r$. Repeating the arguments used in item 2, we obtain the following estimate for j :

$$j = R_b + a > r + R_b.$$

Consequently, there are no roots from Ψ in the j th row. Then there are no roots from Ψ in the j th column as well, a contradiction with $(i, j) \in \Psi$. $\square$

To each block, except for the last one, of the reductive part $\mathfrak{r}$ we put in correspondence a certain minor of the matrix $\mathbb{X}$ in accordance with the following rule. Let $j_1, j_2, \dots, j_{\tilde{r}_k}$ be the numbers of columns in which there are all roots from Ψ that lie to the right of the k th block in $\mathfrak{r}$.

Definition 3.3. The minor in $\mathbb{X}$ that stands at the intersection of $R_{k-1} < i \leq \tilde{r}_k + R_{k-1}$ rows and $j_1, j_2, \dots, j_{\tilde{r}_k}$ columns is called a *principal minor*.

We indicate all principal minors in Example 1 and 2. As above, the minor M_I^J is a minor of the matrix $\mathbb{X}$ with the ordered systems of rows I and columns J .

In Example 1, the minors M_1^2 , $M_{2,3}^{5,6}$, $M_{5,6}^{7,9}$, M_7^8 , $M_{8,9}^{11,12}$, and $M_{11,12}^{13,14}$ are principal. In Example 2, $M_{1,2}^{3,4}$, $M_{3,4}^{5,7}$, M_5^6 , $M_{6,7,8}^{9,10,14}$, $M_{9,10}^{11,12}$, and M_{11}^{12} are principal.

The size of the principal minor that stands to the right of the k th block in the subalgebra $\mathfrak{r}$ is equal to $\tilde{r}_k = \min(r, r_k)$. The number of the principal minors is one less than the number of blocks in $\mathfrak{r}$. Note that no two principal minors have a common column/row.

Lemma 3.4. Let (i, j) be a root from Ψ . Then for any $\tilde{j} < j$ such that $(i, \tilde{j}) \notin \Delta_{\mathfrak{r}}^+$, a root $(\tilde{i}, \tilde{j})$, $\tilde{i} \geq i$, lying in Ψ exists.

PROOF. Let $(i, \tilde{j}) \notin \Delta_{\mathfrak{r}}^+$ and $\tilde{j} < j$. By definition of the base S , there is a root $\xi \in S$ such that $(i, \tilde{j}) - \xi \in \Delta^+$. Lemma 2.1 implies that $\xi = (i, b)$, where $b < \tilde{j}$, or $\xi = (a, \tilde{j})$, where $a > i$. The first case is impossible, because in a row, there is no root from Φ to the right of a root from S ; therefore $\xi = (a, \tilde{j})$, where $a > i$.

For some number l , we have $R_{l-1} < a \leq R_l$. The first assertion of Lemma 3.2 states that $(R_{l-1} + 1, \tilde{j}) \in S \cup \Phi$. We show that this root lies in the system Ψ .

For some k we have $R_{k-1} < \tilde{j} \leq R_k$. Obviously, $(R_k, R_k + 1) \in S$. If $\tilde{j} < R_k$, then $(\tilde{j}, R_k) \in \Delta_{\mathfrak{r}}^+$ and the pair of roots $((a, \tilde{j}), (R_k, R_k + 1))$ is admissible. Consequently, $(\tilde{j}, R_k) \in \Phi$. If $\tilde{j} = R_k$, then $(\tilde{j}, R_k + 1) \in S$. Thus, $(\tilde{j}, R_k + 1) \in S \cup \Phi$ for any $\tilde{j}$ such that $R_{k-1} < \tilde{j} \leq R_k$. Lemma 3.2 and (5) imply that all the roots in the chain

$$(\tilde{j}, R_k + 1), (R_k + 1, R_{k+1} + 1), \dots, (R_{\tilde{s}} + 1, R_{\tilde{s}+1} + 1)$$

lie in Ψ . Next, from Lemma 2.7 it follows that the size of the k th block in $\mathfrak{r}$ is less than r . Then

$$\tilde{j} \leq R_k = R_{k-1} + r_k < R_{k-1} + r.$$

From (5) it follows that the root $(R_{l-1} + 1, \tilde{j})$ is also contained in Ψ .

Thus, we showed that for any $\tilde{j} < j$ there is a number $\tilde{i} = R_{l-1} + 1$ such that $(\tilde{i}, \tilde{j}) \in \Psi$. $\square$

Notation 3.5. Denote by $\mathcal{X}$ the set of matrices x in $\mathfrak{m}$ satisfying the following conditions:

1. The principal minors are triangular,

$$\begin{vmatrix} * & * & \dots & * \\ * & * & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & 0 & \dots & 0 \end{vmatrix},$$

and the numbers on the secondary diagonal are different from zero. In the above diagram we denote by asterisks any numbers from the ground field K .

2. If $(i, j) \in \Psi$ and j is the greatest number for a given i , then $w_{i, \tilde{j}}(x) = 0$ for any $\tilde{j} > j$.
3. Let a principal minor stand at the intersection of the rows $i, i+1, \dots, i+\tilde{r}-1$ and columns $j_1 < j_2 < \dots < j_{\tilde{r}}$. Let

$$(\tilde{i}, \tilde{j}) \notin \Psi, \text{ where } i \leq \tilde{i} < i + \tilde{r} \text{ and } j_{a-1} < \tilde{j} < j_a$$

for some $a \in \{2, \dots, \tilde{r}\}$. Then if the root (i, j_a) stands in the principal minor below the secondary diagonal (in this case $w_{i, j_a}(x) = 0$), then $w_{i, \tilde{j}}(x) = 0$.

We present an example of the set $\mathcal{X} \subset \mathfrak{m}$ in which diagram 1 is a diagram for $\mathfrak{m}$.

$$\begin{pmatrix} 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & * & * & * & * & * & * & * & * & * \\ \hline & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 & 0 & 0 \\ & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline & & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & 0 & 0 & 0 & * & * & 0 & 0 & & & \\ & & 0 & 0 & 0 & * & 0 & 0 & 0 & & & \\ & & 0 & 0 & 0 & * & * & * & * & & & \\ \hline & & & 0 & 0 & & * & * & & & & \\ & & & 0 & 0 & & * & 0 & & & & \\ & & & & & & & & & & 0 & 0 \\ & & & & & & & & & & 0 & 0 \end{pmatrix}$$

Notation 3.6. Denote by T a set of roots $(i, j) \in \Delta^+$ that satisfy the following conditions:

1. There exists a number $\tilde{i} > i$ such that $(\tilde{i}, j) \in \Psi$.
2. In the i th row there are no roots from Ψ .

Denote $\mathcal{T} = \text{Span}\{E_\xi, \xi \in T\}$. Below we give the diagram of Example 1; here the roots from T are denoted by the symbol $\bullet$.




Diagram 4

4. Canonical matrices on N -orbits of general position

First we state two propositions, which will be necessary for the proof of Theorem 1.7.

Remark 4.1. The adjoint action

$$\mathrm{Ad}_{g(t)} x = g(t) \cdot x \cdot g(t)^{-1}$$

on an arbitrary matrix x by the one-parameter subgroup

$$g(t) = E + t \cdot E_\psi \in N, \quad \psi = (u, v) \in \Delta^+$$

reduces to the composition of two transformations:

- 1) the row with number v multiplied by t is added to the row u of the matrix x ;
- 2) the column with the number u multiplied by $-t$ is added to the column with the number v of the matrix x .

Proposition 4.2. *For any $x \in \mathcal{X}$ and any matrix $y \in \mathcal{T}$, there is $g \in N$ such that $\mathrm{Ad}_g(x + y) = x$.*

PROOF. It is sufficient to prove the following assertion. Let j be the number of a column such that all entries of the matrix y in the columns with numbers greater than j are zero. We denote by $\{i_1, \dots, i_p\}$ the numbers of the rows for which the entries of the matrix y in the j th column are equal to zero. We claim that there exists $g \in N$ such that

$$\mathrm{Ad}_g(x + y) = x + \tilde{y},$$

where $\tilde{y} \in \mathcal{T}$ and all the entries of $\tilde{y}$ that lie in the columns with numbers at least j are zero.

Let $i \in \{i_1, \dots, i_p\}$; since $(i, j) \in T$, a root $(\tilde{i}, j) \in \Psi$, $\tilde{i} > i$, exists. Then there exists a principal minor M_I^J , where I is the set of rows and J is the set of columns of the minor, such that $\tilde{i} \in I$ and $j \in J$. Let the principal minor M_I^J stand to the right of the $(c+1)$ th block in $\mathfrak{r}$; denote $k = R_c + 1$ and, as above, $\tilde{r} = \tilde{r}_{c+1} = \min(r, r_{c+1})$. Let

$$I = \{k, k+1, \dots, k+\tilde{r}-1\} \text{ and } J = \{j_1, j_2, \dots, j_{\tilde{r}}\}$$

for some $j_1 < j_2 < \dots < j_{\tilde{r}}$. Denote by l the number for which $j = j_l$. Note that the root $(k+\tilde{r}-l, j_l)$ stands on the secondary diagonal of a principal minor, and thus the value of $w_{k+\tilde{r}-l, j_l}(x)$ does not equal to zero.

Let $i \in \{i_1, \dots, i_p\}$; then $w_{i,j}(y) \neq 0$. Consider the adjoint action on $x + y$ of the element

$$g = \prod_{i=i_1, \dots, i_a} g_i(t_i),$$

where $g_i(t_i) = E + t_i \cdot E_{i, k+\tilde{r}-l}$ and $t_i = -\frac{w_{i,j}(y)}{w_{k+\tilde{r}-l,j}(x)}$.

Then the columns in the matrices $\text{Ad}_g(x + y)$ and x the numbers of which are at least j coincide. Therefore, the entries of the matrix

$$\tilde{y} = \text{Ad}_g(x + y) - x,$$

that lie in the columns with numbers at least j are zero. Hence we obtain $\text{Ad}_g(x + y) = x + \tilde{y}$.

To complete the proof, it remains to prove that $\tilde{y} \in \mathcal{T}$. We show that if $w_{a,b}(\tilde{y}) \neq 0$, then $(a, b) \in T$.

Thus, let $w_{a,b}(\tilde{y}) \neq 0$ and $w_{a,b}(\tilde{y}) \neq w_{a,b}(y)$ for a certain root (a, b) . By Remark 4.1, either $a = i$ or $b = k + \tilde{r} - l$.

1. Let $w_{i,b}(\tilde{y}) \neq 0$ and $w_{i,b}(\tilde{y}) \neq w_{i,b}(y)$, where $b \in \{j_1, j_1 + 1, \dots, j\}$. We show that $(i, b) \in T$. Lemma 3.4 implies that for every number $b < j$, a root $(i_b, b) \in \Psi$ with number $i_b > i$ exists. This, together with the fact that there are no roots from Ψ in the i th row, implies that the root (i, b) lies in T .
2. Let $w_{a, k+\tilde{r}-l}(\tilde{y}) \neq 0$ and $w_{a, k+\tilde{r}-l}(\tilde{y}) \neq w_{a, k+\tilde{r}-l}(y)$ for some a . we show that $(a, k + \tilde{r} - l) \in T$.

- (a) We prove that there are no roots from Ψ in the row with number a . To this end, we first prove that there are no roots from Ψ in the i th column. Assume the contrary. Then the i th row would also contain roots from Ψ , which contradicts the definition of the root system T . Consequently, in the column with number i there are no roots from Ψ . Next, since $w_{a, k+\tilde{r}-l}(y) \neq w_{a, k+\tilde{r}-l}(\tilde{y})$, it follows that the entry $(a, k + \tilde{r} - l)$ of the matrix $x + y$ is changed under the action of g , and thus $w_{a,i}(x + y) \neq 0$. Now assume that in the a th row there is a root (a, v) from Ψ , and let v be the greatest possible. Then, since in the matrices from $\mathcal{X}$ and $\mathcal{T}$ there are zeros to the right of the principal minors, we obtain $i \leq v$. by Lemma 3.4, there is a root from Ψ in the i th column, a contradiction. Therefore, there are no roots from Ψ in the a th row.

(b) Now we show that there is a number $\tilde{a} > a$ such that $(\tilde{a}, k + \tilde{r} - l) \in \Psi$. Assume that there are no roots from Ψ in the $(k + \tilde{r} - l)$ th column. Then there are no roots from S in the $(k + \tilde{r} - l)$ th column. Otherwise Lemma 3.2 and the fact that the row with number $(k + \tilde{r} - l)$ contains roots from the system Ψ imply that there is a root $(\tilde{a}, k + \tilde{r} - l) \in \Psi$ for some $\tilde{a}$. But in the $(k + \tilde{r} - l)$ th column there are no roots from S only if for every $\tilde{c} \leq c$ the size of the $\tilde{c}$ th block in $\mathfrak{r}$ is less than the size of the last block. In particular, the size of the block to the right of which the root $(a, k + \tilde{r} - l)$ stands is less than r . Then the row with number a is a row of a certain principal minor, and since $w_{a,i}(x + y) \neq 0$, there is a root from Ψ in the a th row, which contradicts item (a). Therefore, the $(k + \tilde{r} - l)$ th column contains a root $(\tilde{a}, k + \tilde{r} - l) \in \Psi$.

Now assume that $\tilde{a} < a$. The root $(\tilde{a}, k + \tilde{r} - l)$ lies in Ψ and $i < k + \tilde{r} - l$. Since $(a, i) \notin \Delta_{\mathfrak{r}}^+$, we have $(\tilde{a}, i) \notin \Delta_{\mathfrak{r}}^+$. Then the conditions of Lemma 3.4 are fulfilled, whence it follows that there is a root from Ψ in the i th column, but this contradicts item (a). Consequently, $\tilde{a} > a$.

Items (a) and (b) imply that the root $(a, k + \tilde{r} - l) \in T$.

Thus, all the roots of the form (a, b) , where $w_{a,b}(\tilde{y}) \neq 0$, lie in the root system T . $\square$

Proposition 4.3. *Assume that a parabolic subalgebra $\mathfrak{p}$ is such that there is a root from S in the last column. Then there exists a nonempty Zariski-open subset U in $\mathcal{Y}$ such that the N -orbit of any matrix from U has a nonzero intersection with $\mathcal{X}$.*

PROOF. Consider the linear span

$$\mathcal{Y} + \sum K \cdot E_{\eta}, \quad (7)$$

where summation proceeds over all roots $\eta = (i, j) \in M \setminus \Psi$ such that $i \in I$, $j \in J$, and M_I^J is a principal minor (i.e., the root η lies inside a principal minor).

First we show that there is a nonempty Zariski-open subset V in $\mathcal{Y}$ such that for any $x \in V$ there is a $h \in N$ such that $\text{Ad}_h x$ lies in (7) and any principal minor in $\text{Ad}_h x$ is different from zero.

Let $x \in \mathcal{Y}$. As above, we denote $R_{k-1} = \sum_{t=1}^{k-1} r_t$ and $\tilde{r} = \tilde{r}_k = \min(r, r_k)$.

Let the principal minor M_I^J that stands to the right of the k th block in $\mathfrak{r}$ be

situated at the intersection of $I = \{R_{k-1} + 1, R_{k-1} + 2, \dots, R_{k-1} + \tilde{r}\}$ rows and $J = \{j_1, j_2, \dots, j_{\tilde{r}}\}$ columns.

By Lemma 3.2, there is a root from Ψ in every column of the principal minor, but not in any row of the principal minor there is a root from Ψ . The roots (4, 7) and (4, 8) in Diagram 5 serve as an example. These roots do not lie in the system Ψ , and the fourth row belongs to the set of rows of the principal minors.

Example 3. The diagram for a parabolic subalgebra with the following sizes of diagonal block: (1, 1, 4, 2).




Diagram 5

Let $i \in I$, $j \in J$, and $(i, j) \notin \Psi$. Then $(i, j) \notin S \cup \Phi$. Since j is a column of the principal minor, there is a root from Ψ in the j th column. Therefore, a root $(i, j) \in S$ exists; suppose $\tilde{i} > i$. The adjoint action on x of the element

$$\tilde{h} = E + t \cdot E_{i, \tilde{i}}, \quad t \neq 0, \quad (8)$$

enables one to obtain an arbitrary nonzero number at the entry (i, j) of the matrix x ($w_{\tilde{i}, j}(x) \neq 0$ by the definition of $\mathcal{Y}$). We show that the adjoint action (8) does not change the entries of the matrix x , except for (i, j) . On the one hand, the number that stands at the $(\tilde{i}, j)$ entry of the matrix x is a unique nonzero in the $\tilde{i}$ th row. Indeed, if $w_{\tilde{i}, j}(x) \neq 0$ for some $\tilde{j}$, then $(\tilde{i}, \tilde{j}) \in \Phi$ and $\tilde{j} < j$. But in this case, a root $(b, \tilde{i}) \in S$ exists. Then, by the corollary of Lemma 2.3, there is a root $(a, i) \in S$ in the i th column. Consequently, the pair of roots $((a, i), (\tilde{i}, j))$ is admissible, whence $(i, j) \in \Phi$, which is impossible. On the other hand, to obtain some changes in x upon adjoint action (8), it is necessary that the i th column contain a root from the base S . Again we have a contradiction with $(i, j) \notin \Phi$. Thus we showed that adjoint action (8) does not change the matrix x , except for the entry (i, j) .

Since the roots from S occupy places not above the secondary diagonal in a principal minor, there exists $h \in N$ such that $\text{Ad}_h x$ lies in (7) and

$w_{i,j}(\text{Ad}_h x) \neq 0$ for any root (i, j) that stands not below the secondary diagonal in a principal minor. then the set of $x \in \mathcal{Y}$ for which all principal minors in $\text{Ad}_h x$ are not equal to zero is a nonempty Zariski-open set; denote it by V .

Now we turn to a proof of the proposition. Denote $x' = \text{Ad}_h x$, and let x' lie in (7). Suppose $x \in V$. Then all principal minors in x' are different from zero.

Consider the first r_1 rows in x' . Obviously, they satisfy conditions 1–3 imposed on the matrices from $\mathcal{X}$, because they do not contain roots from Φ and in any row (column) there is no more one root from S .

Now assume that for any row the number of which is no greater than R_{k-1} , the matrix x' satisfies requirements 1–3 in the definition of the set $\mathcal{X}$. To prove the proposition, it is sufficient to show that there exists a nonempty Zariski-open subset $U \subset \mathcal{Y}$ such that for any $x \in U$ there is $g \in N$ such that the matrix $\text{Ad}_g x' = \text{Ad}_{gh} x$ satisfies requirements 1–3 in the definition of the set $\mathcal{X}$ for rows the numbers of which do not exceed R_k . We represent the part of the diagram that corresponds to the rows $R_{k-1} + 1, R_{k-1} + 2, \dots, R_k$. Lemma 3.2 implies that it has the following form

	j_1	j_2	$\dots$	$j_{\widetilde{r}}$					
1	$\boxtimes$	$\dots$	$\boxtimes$	$\dots$	$\boxtimes$	$\dots$	$\times$		$R_{k-1} + 1$
$\dots$	$\dots\d$								

Let the principal minor M_I^J stand to the right of the block with number k in $\mathfrak{r}$ at the intersection of the rows $I = \{R_{k-1} + 1, R_{k-1} + 2, \dots, R_{k-1} + \tilde{r}\}$ and the columns $J = \{j_1, j_2, \dots, j_{\tilde{r}}\}$, where $j_1 < j_2 < \dots < j_{\tilde{r}}$. Denote by U' the nonempty Zariski-open subset of those $x \in \mathcal{Y}$ for which the principal minor M_I^J in x' is different from zero. We show that for any $x \in U = U' \cap V$, there is $g \in N$ such that the matrix $\text{Ad}_g x'$ satisfies the requirements of the definition of the set $\mathcal{X}$ for rows the numbers of which does not exceed R_k .

Thus, let $x \in U$; therefore the principal minor M_I^J is different from zero. Then we can put it in triangular form with the help of the adjoint action of elements from N of the form

$$\tilde{g}_a = E + t_1^a \cdot E_{j_1, j_a} + \dots + t_{a-1}^a \cdot E_{j_{a-1}, j_a}, \quad a = 2, \dots, \tilde{r}, \quad (9)$$

choosing appropriate values of t_i^a , $i = 1, \dots, a - 1$. In such a way we achieve the fulfillment of condition 1 in the definition of $\mathcal{X}$ for rows the number of which do not exceed R_k .

Next, let $(i, j) \in M \setminus \Psi$, where $i \in I$ and $j > j_{\tilde{r}}$, moreover $w_{i,j}(x') \neq 0$. Similarly, we act on x' by conjugation by the element

$$\widehat{g}_j = E + t_1 \cdot E_{j_1,j} + \dots + t_{\tilde{r}} \cdot E_{j_{\tilde{r}},j}, \quad (10)$$

where $t_1, \dots, t_{\tilde{r}}$ are appropriate values. We achieve that the entry (i, j) of the matrix $\text{Ad}_{\widehat{g}_j} x'$ becomes zero and the conjugation by $\widehat{g}_j$ does not change the other entries of the matrix in the $R_{k-1} + 1, R_{k-1} + 2, \dots, R_k$ rows. The latter means that condition 2 in the definition of the set $\mathcal{X}$ is satisfied for the rows the numbers of which do not exceed R_k .

Now we assume that the principal minor to the right of the k th block is triangular. Let $w_{i,j}(x') \neq 0$ for a root (i, j) , where $i \in I$, $j_{a-1} < j < j_a$ for some $j_{a-1}, j_a \in J$, and the root (i, j_a) lies below the secondary diagonal in the principal minor. It is sufficient to consider the conjugation by element (9), replacing j by j_a . Thus, the condition 3 in the definition of $\mathcal{X}$ for rows the numbers of which do not exceed R_k is satisfied. $\square$

Now let $\mathcal{S}$ be the set of denominators generated by minors M_ξ , $\xi \in S$. We form the localization $K[\mathfrak{m}]_{\mathcal{S}}$ of the algebra $K[\mathfrak{m}]$ on $\mathcal{S}$.

Consider the Zariski-open subset in $\mathfrak{m}$

$$U_0 = \{x \in \mathfrak{m} : M_\xi(x) \neq 0, \text{ для любого } \xi \in S\}.$$

Then $K[\mathfrak{m}]_{\mathcal{S}} = K[U_0]$. We define a regular map

$$\pi : U_0 \rightarrow \mathcal{Y} \quad (11)$$

by the rule

$$x \in U_0 \mapsto \sum_{\xi \in S} \frac{M_\xi(x)}{\widehat{M}_\xi(x)} \cdot E_\xi + \sum_{\varphi \in \Phi} \frac{L_\varphi(x)}{M_{\xi_1}(x) \widehat{M}_{\xi_2}(x)} \cdot E_\varphi \in \mathcal{Y},$$

where

- 1) $\widehat{M}_\gamma(x)$ is a number $\prod M_{\gamma'}(x)$, and the product proceeds over all roots γ' such that $\gamma' < \gamma$ and γ' is the greatest root in the sense of the order $<$ introduced above;
- 2) the root $\varphi \in \Phi$ from the second sum corresponds to the admissible pair $q = (\xi_1, \xi_2)$.

Then $M_\xi(x) = M_\xi(\pi(x))$, $L_\varphi(x) = L_\varphi(\pi(x))$ for any $\xi \in S$, $\varphi \in \Phi$. Since M_ξ and L_φ are invariants, we have $\pi(\text{Ad}_g x) = \pi(x)$ for any $g \in N$.

Now we are in a position to prove theorem 1.7.

PROOF OF THEOREM 1.7 proceeds by induction on n .

For $n = 2$ the statement is obvious. Assume that the theorem is valid for matrices of sizes lesser than n . We prove it for matrices of size n . Let the reductive part $\mathfrak{r}$ of a parabolic subalgebra $\mathfrak{p}$ consists of blocks of sizes $(r_1, r_2, \dots, r_s)$, where $r_1 + r_2 + \dots + r_s = n$. As above, $r = r_s$. Introduce the notation

$$\mathfrak{z}_0 = \left\{ \left(\begin{array}{ccc|c} & & & * \\ & & & \dots \\ & & & * \\ \hline 0 & \dots & 0 & 0 \\ & \dots & & \dots \\ 0 & \dots & 0 & 0 \end{array} \right) \right\}^r, \quad \mathcal{Z}_0 = \exp \mathfrak{z}_0.$$

Since $\mathfrak{z}_0$ is an ideal in the subalgebra $\mathfrak{n}$, we have $\text{Ad}_N \mathfrak{z}_0 = \mathfrak{z}_0$. We identify

$$\mathfrak{gl}(n-1, K) \text{ with the subalgebra } \left\{ \left(\begin{array}{ccc|c} * & \dots & * & 0 \\ & \dots & & \dots \\ * & \dots & * & 0 \\ \hline 0 & \dots & 0 & 0 \end{array} \right) \right\} \text{ in } \mathfrak{gl}(n, K)$$

and

$$\text{GL}(n-1, K) \text{ with the subgroup } \left\{ \left(\begin{array}{ccc|c} * & \dots & * & 0 \\ & \dots & & \dots \\ * & \dots & * & 0 \\ \hline 0 & \dots & 0 & 1 \end{array} \right) \right\} \text{ in } \text{GL}(n, K).$$

Let N_1 be the subgroup of upper triangular matrices with unities on the principal diagonal in $\text{GL}(n-1, K)$, $\mathfrak{n}_1 = \text{Lie } N_1$, $\mathfrak{m}_1 = \mathfrak{m} \cap \mathfrak{n}_1$, and let $\mathcal{Y}_1$ be the natural projection of $\mathcal{Y}$ to $\mathfrak{m}_1$.

Any element $x \in \mathfrak{m}$ is uniquely represented in the form $x = x_1 + z_1$, where $x_1 \in \mathfrak{m}_1$ and $z_1 \in \mathfrak{z}_0$. By the inductive assumption, there exists a nonempty Zariski-open subset $U_1 \subset \mathfrak{m}_1$ such that for any $x_1 \in U_1$ there is $g' \in N_1$ such that $\text{Ad}_{g'} x_1 \in \mathcal{Y}_1$. Since $\mathfrak{m} = \mathfrak{m}_1 \oplus \mathfrak{z}_0$, it follows that $U_1 + \mathfrak{z}_0$ is an open subset in $\mathfrak{m}$.

Proposition 4.3 implies the existence of a Zariski-open subset V in $\mathcal{Y}$ such that for any $y \in V$, there is an element $g_1 \in N$ such that $\tilde{y} = \text{Ad}_{g_1} y \in \mathcal{X}$. The preimage $\pi^{-1}(V)$ of the set V under map (11) is a Zariski-open subset in $\mathfrak{m}$.

Denote

$$U = \begin{cases} U_1 + \mathfrak{z}_0 & \text{if there are no roots from } S \text{ in the } n\text{th column;} \\ (U_1 + \mathfrak{z}_0) \cap \pi^{-1}(V) & \text{if there is a root from } S \text{ in the } n\text{th column.} \end{cases}$$

From what we have said above, U is a Zariski-open subset in $\mathfrak{m}$.

PART I. We show that for any $x \in U$ there exists $g \in N$ such that $\text{Ad}_g x \in \mathcal{Y}$.

Thus, let $x \in U$. Denote $x' = \text{Ad}_{g'} x$, $y_0 = \text{Ad}_{g'} x_1$, $z_0 = \text{Ad}_{g'} z_1$. We obtain

$$x' = \text{Ad}_{g'} x = \text{Ad}_{g'} x_1 + \text{Ad}_{g'} z_1 = y_0 + z_0,$$

где $y_0 \in \mathcal{Y}_1$, $z_0 \in \mathfrak{z}_0$.

1. Consider the case where there are no roots from the base S in the n th column. In this case, $U = U_1 + \mathfrak{z}_0$. We show that there is $g \in N$ such that $\text{Ad}_g x = y_0$. Lemma 2.1 implies that for any number i , where $1 \leq i \leq n-r$, there is a root $\xi \in S$ lying in the i th row (i.e., $\xi = (i, a_i)$ for some number a_i). Since $y_0 \in \mathcal{Y}_1$, we have $w_{i,a_i}(x') = w_{i,a_i}(y_0) \neq 0$. Denote

$$\begin{aligned} g_{n-r} &= E + t_{n-r} \cdot E_{a_{n-r},n} \in \mathcal{Z}_0, \quad \text{where} \quad t_{n-r} = \frac{w_{n-r,n}(z_0)}{w_{n-r,a_{n-r}}(y_0)}; \\ g_i &= E + t_i \cdot E_{a_i,n} \in \mathcal{Z}_0, \quad \text{where} \quad t_i = \frac{w_{i,n}(\text{Ad}_{g_{i+1} \dots g_{n-r}} z_0)}{w_{i,a_i}(y_0)}. \end{aligned} \tag{12}$$

Then

$$w_{a,n}(\text{Ad}_{g_i g_{i+1} \dots g_{n-r}} x') = 0$$

for any $a \geq i$. Denote

$$g = g_1 \cdot g_2 \cdot \dots \cdot g_{n-r} \cdot g'.$$

We obtain

$$\begin{aligned} \text{Ad}_g x &= \text{Ad}_{g_1 g_2 \dots g_{n-r}} \text{Ad}_{g'} x = \text{Ad}_{g_1 g_2 \dots g_{n-r}} x' = \\ &= \text{Ad}_{g_1 g_2 \dots g_{n-r}} (y_0 + z_0) = y_0 \in \mathcal{Y}, \end{aligned}$$

which proves the statement in case 1.

2. Now consider the case where a root from S lies in the n th column. Here $U = (U_1 + \mathfrak{z}_0) \cap \pi^{-1}(V)$. Denote by $(\tilde{m}, n)$ the root from S lying in the n th column. Let $(m, n) \in S \cup \Phi$ be a root which has the least number m .

Lemma 2.1 implies that for every $(i, n) \in M$, where $i > \tilde{m}$, a root from S exists in the i th row. Similarly to the first item, there is an element $h \in \mathcal{Z}_0$ such that $\text{Ad}_h x'$ has zeros in the last column below the root $(\tilde{m}, n)$ (i.e., $w_{a,n}(\text{Ad}_h x') = 0$ for any $a > \tilde{m}$).

Now let $(i, n) \notin S \cup \Phi$ and $m < i < \tilde{m}$. Similarly to the first part of the proof of Proposition 4.3, acting on x' by conjugation by element (8) for $\tilde{i} = \tilde{m}$, we can obtain

$$w_{i,n}(\text{Ad}_h x') = 0 \text{ and } w_{a,b}(\text{Ad}_h x') = w_{a,b}(x')$$

for any root $(a, b) \neq (i, n)$.

In view of the above-said, in the sequel we assume that for x' , $w_{i,n}(x') = 0$ for any root $(i, n) \notin S \cup \Phi$ such that $i > m$.

Consider the subspace $\mathfrak{z} \subset \mathfrak{z}_0$ that consists of $z \in \mathfrak{z}_0$ for which $w_{i,n}(z) = 0$ for $i \geq m$. Denote by z the natural projection of z_0 to $\mathfrak{z}$. Then the matrix $y = x' - z$ lies in $\mathcal{Y}$. Therefore, $x' = y + z$, $y \in \mathcal{Y}$, $z \in \mathfrak{z}$. From the definition of invariants, $y = \pi(x')$. Since $x' \in \pi^{-1}(V)$, we have $y \in V$.

From Proposition 4.3 it follows that there exists an element $g_1 \in N$ such that $\tilde{y} = \text{Ad}_{g_1} y \in \mathcal{X}$. Then $\text{Ad}_{g_1} x' = \tilde{y} + \text{Ad}_{g_1} z$. Since $\mathfrak{z}$ is an ideal of $\mathfrak{n}$, we have $\text{Ad}_{g_1} z \in \mathfrak{z}$.

We show that there exists $g_2 \in \mathcal{Z}_0$ such that

$$\text{Ad}_{g_2 g_1} x' = \tilde{y} + \tilde{z},$$

where $\tilde{y} \in \mathcal{X}$ and $\tilde{z} \in \mathcal{T} \cap \mathfrak{z}$. Let $\zeta = (i, n) \notin T$, $i < m$. In the n th column, there is a root from S and thus from Ψ . Then, by definition of T , there exists a principal minor M_I^J , where I is a set of numbers of rows and J is the set of numbers of columns, such that $i \in I$. Let $\tilde{r}$ be its size and $J = \{j_1, \dots, j_{\tilde{r}}\}$, $j_1 < \dots < j_{\tilde{r}}$. Since $\tilde{y} \in \mathcal{X}$, the principal minor M_I^J is different from zero. We choose numbers $t_1, \dots, t_{\tilde{r}}$ in such a way that for the element

$$g_\zeta = E + t_1 \cdot E_{j_1, n} + \dots + t_{\tilde{r}} \cdot E_{j_{\tilde{r}}, n}$$

the following conditions are satisfied:

- (a) $\text{Ad}_{g_\zeta}(\tilde{y} + \text{Ad}_{g_1} z) = \tilde{y} + z'$, where $z' \in \mathfrak{z}$;
- (b) $w_{i,n}(\text{Ad}_{g_\zeta g_1} z) = 0$;
- (c) if $i \neq j$, then the adjoint action of g_ζ does not change the entry (j, n) of the matrix $\tilde{y} + \text{Ad}_{g_1} z$.

Then for $g_2 = \prod_{\zeta=(i,n) \notin T} g_\zeta \in \mathcal{Z}_0$, we have

$$\mathrm{Ad}_{g_2 g_1} x' = \tilde{y} + \tilde{z}, \text{ where } \tilde{y} \in \mathcal{X} \text{ and } \tilde{z} \in \mathcal{T} \cap \mathfrak{z}.$$

Next, Proposition 4.2 implies the existence of $g_3 \in N$ such that

$$\mathrm{Ad}_{g_3}(\tilde{y} + \tilde{z}) = \tilde{y}.$$

We obtain

$$\mathrm{Ad}_{g_3 g_2 g_1} x' = \mathrm{Ad}_{g_3 g_2 g_1}(y + z) = \mathrm{Ad}_{g_3}(\tilde{y} + \tilde{z}) = \tilde{y}.$$

It remains to apply the inverse action g_1^{-1} :

$$\mathrm{Ad}_{g_1^{-1} g_3 g_2 g_1 g'} x = \mathrm{Ad}_{g_1^{-1} g_3 g_2 g_1} x' = \mathrm{Ad}_{g_1^{-1}} \tilde{y} = \mathrm{Ad}_{g_1^{-1}}(\mathrm{Ad}_{g_1} y) = y \in \mathcal{Y}.$$

PART II. Uniqueness follows from the fact that the map (11) is constant on an N -orbit. $\square$

Consider the localization $K[\mathfrak{m}]_S^N$ of the algebra of invariants $K[\mathfrak{m}]^N$ with respect to $\mathcal{S}$. Since the minors M_ξ are N -invariants, we have

$$K[\mathfrak{m}]_S^N = (K[\mathfrak{m}]_S)^N.$$

The following theorem was proved in [PS]; we repeat its proof to maintain the understanding of the matter.

Theorem 4.4. *The ring $K[\mathfrak{m}]_S^N$ is the ring of polynomials in $M_\xi^{\pm 1}$, $\xi \in S$, and in L_φ , $\varphi \in \Phi$.*

PROOF. Consider the restriction homomorphism $\varpi : f \mapsto f|_{\mathcal{Y}}$ of the algebra $K[\mathfrak{m}]^N$ to $K[\mathcal{Y}]$. The image $\varpi(M_\xi)$ is equal to the product

$$\pm x_\xi x_{\xi_1} \dots x_{\xi_t},$$

where every ξ_i is less than ξ in the sense of above order $>$. We extend ϖ to a homomorphism

$$\varpi_S : K[\mathfrak{m}]_S^N \rightarrow K[\mathcal{Y}]_S,$$

where $K[\mathcal{Y}]_S$ is a localization of $K[\mathcal{Y}]$ with respect to x_ξ , $\xi \in S$. We show that ϖ_S is an isomorphism.

If $f \in \text{Ker } p_S$, then $f(\text{Ad}_N \mathcal{Y}) = 0$. Since, by Theorem 1.7, $\text{Ad}_N \mathcal{Y}$ contains a Zariski-open subset, we have $f = 0$. Consequently, ϖ_S is an embedding of $K[\mathfrak{m}]_S^N$ in $K[\mathcal{Y}]_S$. Next, we write the formulas in the form

$$\begin{aligned}\varpi(M_\xi) &= \pm x_\xi \varpi(\widehat{M}_\xi), \\ \varpi(L_\varphi) &= \pm x_\varphi \varpi(M_\gamma) \varpi(\widehat{M}_{\gamma'}),\end{aligned}$$

where the root $\varphi \in \Phi$ corresponds to the admissible pair (γ, γ') , and $\widehat{M}_\xi$ ($\widehat{M}_{\gamma'}$, respectively) is the product $\prod M_{\xi_1}$ ($\prod M_{\gamma_1}$, respectively), and the product proceeds over the greatest roots ξ_1 such that ξ_1 is less than ξ (γ' , respectively) in the sense of the order $<$. We get

$$\begin{aligned}\varpi_S(M_\xi \widehat{M}_\xi^{-1}) &= \pm x_\xi, \\ \varpi_S(L_\varphi M_\gamma^{-1} \widehat{M}_{\gamma'}^{-1}) &= \pm x_\varphi.\end{aligned}\tag{13}$$

From (13) it follows that the image of ϖ_S coincides with $K[\mathcal{Y}]_S$. Thus, ϖ_S is an isomorphism. $\square$

We obtain Theorem 1.8 as a consequence of Theorem 4.4.

References

- [B] M. Brion, Représentations exceptionnelles des groupes semi-simples, *Ann. Scient. Ec. Norm. Sup.* **18** (1985), 345–387.
- [GG] M. Goto and F. Grosshans, Semisimple Lie algebras, *Lect. Notes in Pure Appl. Math.*, vol. 38, 1978.
- [K] H. Kraft, Geometrische Methoden in der Invariantentheorie, Friedr. Vieweg and Sohn, Braunschweig/Wiesbaden (1985).
- [PS] A. N. Panov and V. V. Sevostyanova, Regular N -orbits in the nilradical of a parabolic subalgebra, *Vestnik SamGU*, **7**(57) (2007), 152–161. See also <http://arxiv.org/abs/1203.2754>.
- [PV] V. L. Popov and E. B. Vinberg, Invariant theory, in: *Progress in Science and Technology*, VINITI, Moscow (1989), pp. 137–309.
- [R] R. W. Richardson, Conjugacy classes in parabolic subgroups of semisimple algebraic groups, *Bull. London Math. Soc.* **6** (1974), 21–24.

DEPARTMENT OF MECHANICS AND MATHEMATICS, SAMARA STATE UNIVERSITY,
RUSSIA

E-mail address: victoria.sevostyanova@gmail.com